

Transmission time of selected tick-borne disease agents.

Pathogen	Disease	Vector	Transmission Time	Reference
<i>Anaplasma phagocytophilum</i>	Granulocytic anaplasmosis	<i>Ixodes scapularis</i>	4 to 48 hours	Nicholson et al. 2010
		<i>Ixodes pacificus</i>		
		<i>Ixodes scapularis</i>	< 36 hours	Katavolos et al. 1998
		<i>Ixodes scapularis</i>	< 24 hours	des Vignes et al. 2001
<i>Babesia microti</i>	Human babesiosis	<i>Ixodes dammini</i>	40-48 hours	Hodzic et al. 1998
			< 36 to 48 hours	Piesman et al. 1980
			< 48 hours	Crippa et al. 2002
			> 17 hours	Kahl et al. 1998
<i>Borrelia afzelii</i>	Lyme disease-like	<i>Ixodes scapularis</i>	< 24 hours	Piesman et al. 1987
			< 48 to 72 hours	Vignes et al. 2001
			< 48 hours	Meiners et al. 2006
		<i>Ixodes ricinus</i>		
<i>Borrelia burgdorferi</i>	Lyme disease	<i>Amblyomma americanum</i>	36-48 hours	Thomas et al., 2017
		<i>Rhipicephalus sanguineus</i>	4 to 48 hours	Nicholson et al. 2010
		<i>Dermacentor variabilis</i>		
<i>Ehrlichia canis</i>	Canine ehrlichiosis	<i>Rhipicephalus sanguineus</i>	3 hours	Fourie et al. 2013
		<i>Dermacentor variabilis</i>		
		<i>Dermacentor andersoni</i>	4 to 48 hours	Nicholson et al. 2010
<i>Rickettsia rickettsii</i>	Rocky Mountain spotted fever	<i>Rhipicephalus sanguineus</i>		
		<i>Amblyomma aureolatum</i>	> 10 hours	Saraiva et al. 2014
<i>Theileria parva</i>	Theileriosis; East Coast fever	<i>Rhipicephalus appendiculatus</i>	< 24 to 72 hours	Ochanda et al. 1988
<i>Powassan virus</i>	Powassan virus disease	<i>Ixodes scapularis</i>	< 15 minutes	Ebel et al. 2004

Transmission time of selected tick-borne disease agents.

References:

- Crippa M, O Rais, L Gern. 2002. Investigations on the mode and dynamics of transmission and infectivity of *Borrelia burgdorferi* sensu stricto and *Borrelia afzelii* in *Ixodes ricinus* ticks. Vector Borne Zoonotic Dis 2: 3–9
- des Vignes, F, J Piesman, R Heffernan, TL Schulze, KC Stafford, D Fish. 2001. Effect of tick removal on transmission of *Borrelia burgdorferi* and *Ehrlichia phagocytophila* by *Ixodes scapularis* nymphs. J Infect Dis 183: 773–778
- Ebel GD, LD Kramer. 2004. Short report: duration of tick attachment required for transmission of *Powassan* virus by deer ticks. Am J Trop Med Hyg 71: 268–271
- Fourie, JJ, D Stanneck, HG Luus, F Beugnet, M Wijnveld, F Jongejan. 2013. Transmission of *Ehrlichia canis* by *Rhipicephalus sanguineus* ticks feeding on dogs and artificial membranes. Veterinary Parasitology 197: 595–603
- Hodzic E, Fish D, Maretzki CM, et al. 1998. Acquisition and transmission of the agent of human granulocytic ehrlichiosis by *Ixodes scapularis* ticks. J Clin Microbiol. 36: 3574–8.
- Kahl, O, C Janetzki-Mittmann, JS Gray, et al. 1998. Risk of infection with *Borrelia burgdorferi* sensu lato for a host in relation to the duration of nymphal *Ixodes ricinus* feeding and the method of tick removal. J Am Vet Med Assoc 287: 41–52
- Katavolos, P, PM Armstrong, JE Dawson, et al. 1998. Duration of tick attachment required for transmission of granulocytic ehrlichiosis. J. Infect. Dis. 177: 1422–1425
- Meiners T, B Hammer, UB Gobel, et al. 2006 Determining the tick scutal index allow assessment of tick feeding duration and estimation of infection risk with *Borrelia burgdorferi* sensu lato in person bitten by an *Ixodes ricinus* nymph. Int J Med Microbiol 40: 103–107
- Nicholson, WL, KE Allen, JH McQuiston, EB Breitschwerdt, and SE Little. 2010. The increasing recognition of rickettsial pathogens in dogs and people. Trends in Parasitology 26: 205–212
- Ochanda H, AS Young, JJ Mutugi, et al. 1988. The effect of temperature on the rate of transmission of *Theileria parva parva* infection to cattle by its tick vector, *Rhipicephalus appendiculatus*. Parasitology 97: 239–245
- Piesman J, A Spielman. 1980. Human babesiosis on Nantucket Island: prevalence of *Babesia microti* in ticks. Am J Trop Med Hyg 29: 742–746
- Piesman J, TN Mather, RJ Sinsky, et al. 1987. Duration of tick attachment and *Borrelia burgdorferi* transmission. J Clin Microbiol 25: 557–558
- Piesman J, GO Maupin, EG Campos, et al., 1991. Duration of adult female *Ixodes dammini* attachment and transmission of *Borrelia burgdorferi*, with description of a needle aspiration isolation method. J Infect Dis 163: 895–897
- Saraiva DG, Soares HS, Soares JF, et al., 2014. Feeding period required by *Amblyomma aureolatum* ticks for transmission of *Rickettsia rickettsii* to vertebrate hosts. Emerg Infect Dis. 20: 1504–10
- Thomas JW, Ohmes CM, Payton ME, et al., 2017. Minimum transmission time of *Cytauxzoon felis* by *Amblyomma americanum* to domestic cats in relation to duration of infestation, and investigation of ingestion of infected ticks as a potential route of transmission. J Fel Med Surg. PMID: 28151047.